arXiv:1212.0870v1 [stat.ME] 4 Dec 2012

Errors-in-variables beta regression models

Jalmar M. F. Carrasco

Departamento de Estatística, Universidade de São Paulo, Brazil

Silvia L. P. Ferrari [1]

Departamento de Estatística, Universidade de São Paulo, Brazil

Reinaldo B. Arellano-Valle

Departamento de Estadística, Pontifícia Universidad Católica de Chile, Chile

---

Abstract

Beta regression models provide an adequate approach for modeling continuous outcomes limited to the interval $(0, 1)$. This paper deals with an extension of beta regression models that allows for explanatory variables to be measured with error. We follow the structural approach, in which the covariates measured with error are assumed to be random variables. We present three estimation methods, namely maximum likelihood, maximum pseudo-likelihood and regression calibration. Monte Carlo simulations are used to evaluate the performance of the proposed estimators and the naïve estimator. Also, a residual analysis for beta regression models with measurement errors is proposed. Our results are illustrated in a real data set.

Keywords: Beta regression model; Errors-in-variables model; Gauss-Hermite quadrature; Maximum likelihood; Maximum pseudo-likelihood; Regression calibration.

---

1. Introduction

Errors-in-variables models, also called measurement error models, are widely applicable in many research areas since they allow for the presence of explicative

---

[1] Corresponding author: Departamento de Estatística, Universidade de São Paulo, Rua do Matão, 1010, 05508-090, São Paulo, SP, Brazil. e-mail:silviaferrari.usp@gmail.com

variables that are measured with errors or that cannot be observed directly (latent variables). Many examples and applications of these models are considered in the books by Fuller (1987), Carroll *et al.* (2006) and Cheng and Van Ness (1999). It is well known that measurement errors cause biased and inconsistent parameter estimates and lead to erroneous conclusions in inferential analysis. Errors-in-variables models are specified in such a way that the distribution of the response variable, y, is assumed to depend on covariates, x, which are imprecisely measured, and observable variables, w, are seen as surrogates for the unobservable true covariates. The classical linear errors-in-variables model has been extensively discussed in the literature, particularly under the normality assumption for the distribution of the unobservable variables. For a systematic review of such models see Fuller (1987) and Cheng and Van Ness (1999); see also Arellano-Valle and Bolfarine (1996) and Castro *et al.* (2008). .

The beta regression models provide an adequate approach for modeling continuous outcomes limited to the interval $(0, 1)$, or more generally, limited to any open interval (a, b) as long as the limits are known (Ferrari and Cribari-Neto, 2004). Although the literature on beta regression has grown fast in the last few years, errors-in-variables models with beta distributed outcomes is an unexplored area.

A beta regression model assumes that the response variable, y, has a beta distribution with probability density function

$$f(y; \mu, \phi) = \frac{\Gamma(\phi)}{\Gamma(\mu\phi)\Gamma[(1-\mu)\phi]} y^{\mu\phi-1}(1-y)^{(1-\mu)\phi-1}, \;\; 0 < y < 1, \tag{1}$$

where $\Gamma(\cdot)$ is the gamma function, $0 < \mu < 1$ and $\phi > 0$, and we write $y \sim \mathrm{Beta}(\mu, \phi)$. Here, $\mu = \mathrm{E}(y)$ and ϕ is regarded as a precision parameter since $\mathrm{Var}(y) = \mu(1-\mu)/(1+\phi)$. For independent observations $y_1, y_2, \ldots, y_n$, where each y_t follows a beta density (1) with mean μ_t and unknown precision parameter ϕ, the beta regression model defined by Ferrari and Cribari-Neto (2004) assumes that

$$g(\mu_t) = \mathbf{z}_t^\top \boldsymbol{\alpha}, \tag{2}$$

with $\boldsymbol{\alpha} \in \mathbb{R}^{p_\alpha}$ being a column vector of unknown parameters, and with $\mathbf{z}_t^\top = (z_{t1}, \ldots, z_{tp_\alpha})$ being a vector of p_α fixed covariates $(p_\alpha < n)$. The link function $g(\cdot) : (0, 1) \to \mathbb{R}$ is assumed to be a continuous, strictly monotone and twice differentiable function. There are many possible choices for $g(\cdot)$, for instance, the logit link, $g(\mu_t) = \log[\mu_t/(1-\mu_t)]$, the probit link, $g(\mu_t) = \Phi^{-1}(\mu_t)$, where $\Phi(\cdot)$ is the cumulative distribution function of the standard normal distribution, and the complementary log-log link, $g(\mu_t) = \log[-\log(1-\mu_t)]$.

Extensions for the beta regression model proposed by Ferrari and Cribari-Neto (2004) that allow the precision parameter to vary across observations, or that involve non-linear structures for the regression specification of the mean and the precision parameter, are presented by Smithson and Verkuilen (2006), Simas *et al.* (2010), among others. The beta regression model with linear specification for the transformed mean and precision parameter is given by (1), (2) and

$$h(\phi_t) \;=\; \mathbf{v}_t^\top \boldsymbol{\gamma}, \tag{3}$$

where $\boldsymbol{\gamma} \in \mathbb{R}^{p_\gamma}$ ($p_\alpha + p_\gamma < n$) is a column vector of unknown parameters, $\mathbf{v}_t = (v_{t1}, \cdots, v_{tp_\gamma})^\top$ is a vector of fixed covariates, $h(\cdot) : (0,\infty) \longrightarrow \mathbb{R}$ is a strictly monotone, twice differentiable link function. A possible choice for $h(\cdot)$ is $h(\phi_t) = \log(\phi_t)$.

The purpose of this paper is to extend the beta regression model (1)-(3) to the situation where some covariates are not directly measured or are measured with error. A practical application of errors-in-variables beta regression models will be illustrated in a study of the risk of coronary heart disease as a function of low-density lipoprotein (LDL) cholesterol level ("bad cholesterol") and body mass index (BMI). The dataset consists of observations of systolic blood pressure (SBP), diastolic blood pressure (DBP), BMI and total cholesterol level (TC) in a group of 182 smoking women aged 50 to 87 years. The total cholesterol may be considered as a surrogate of LDL, which is a covariate of interest, and whose direct measure is more expensive and time consuming. The difference between SBP and DBP results in what is known as the pulse pressure, $PP = SBP - DBP$, and the relative pulse pressure is $RPP = (SBP - DBP)/SBP = PP/SBP$. Small values of RPP, $RPP < 25\%$ say, is indicative of risk of heart disease (American College of Surgeons, 2008, p. 58). Notice that the response variable, RPP, is continuous and limited to the unit interval, and that one of the covariates, namely LDL, is not measured directly.

This paper is organized as follows. In Section 2, we present an errors-in-variables beta regression model under the structural approach, and the corresponding likelihood function. In Section 3, we present three different estimation methods, namely maximum likelihood, maximum pseudo-likelihood, and regression calibration. In Section 4, we perform a simulation study to evaluate and compare the performance of the three estimation approaches. In Section 5, we propose a residual analysis. Section 6 presents an application of the proposed model. Concluding remarks are presented in Section 7.

2. Model and likelihood

Let $y_1, \ldots, y_n$ be independent observable random variables arising from a sample of size n, such that y_t has a beta distribution with probability density function (1) with parameters $\mu = \mu_t$ and $\phi = \phi_t$. In the following, we assume that μ_t and ϕ_t may depend on covariates and unknown parameters. In practice, some covariates may not be precisely observed, but, instead, may be obtained with error. The model considered in this paper assumes a linear structure for the specification of the mean and the precision parameters, and also assumes that both specifications may involve covariates measured with error. Specifically, we replace the mean submodel (2) and the precision submodel (3) by

$$\begin{aligned} g(\mu_t) &= \mathbf{z}_t^\top \boldsymbol{\alpha} + \mathbf{x}_t^\top \boldsymbol{\beta}, && (4)\\ h(\phi_t) &= \mathbf{v}_t^\top \boldsymbol{\gamma} + \mathbf{m}_t^\top \boldsymbol{\lambda}, && (5) \end{aligned}$$

respectively, where $\boldsymbol{\beta} \in \mathbb{R}^{p_\beta}$, $\boldsymbol{\lambda} \in \mathbb{R}^{p_\lambda}$ are column vectors of unknown parameters, $\mathbf{x}_t = (x_{t1}, \cdots, x_{tp_\beta})^\top$ and $\mathbf{m}_t = (m_{t1}, \cdots, m_{tp_\lambda})^\top$ $(p_\alpha + p_\beta + p_\gamma + p_\lambda < n)$ are unobservable (latent) covariates, in the sense that they are observed with error. The vectors of covariates measured without error, $\mathbf{z}_t$ and $\mathbf{v}_t$, may contain variables in common, and likewise, $\mathbf{x}_t$ and $\mathbf{m}_t$. Let $\mathbf{s}_t$ be the vector containing all the unobservable covariates. For $t = 1, \ldots, n$, the random vector $\mathbf{w}_t$ is observed in place of $\mathbf{s}_t$, and it is assumed that

$$\mathbf{w}_t = \boldsymbol{\tau}_0 + \boldsymbol{\tau}_1 \circ \mathbf{s}_t + \mathbf{e}_t, \qquad (6)$$

where $\mathbf{e}_t$ is a vector of random errors, $\boldsymbol{\tau}_0$ and $\boldsymbol{\tau}_1$ are (possibly unknown) parameter vectors and $\circ$ represents the Hadamard (elementwise) product. The parameter vectors $\boldsymbol{\tau}_0$ and $\boldsymbol{\tau}_1$ can be interpreted as the additive and multiplicative biases of the measurement error mechanism, respectively. If $\boldsymbol{\tau}_0$ is a vector of zeros and $\boldsymbol{\tau}_1$ is a vector of ones, we have the classical additive model $\mathbf{w}_t = \mathbf{s}_t + \mathbf{e}_t$. Here, we follow the structural approach, in which the unobservable covariates are regarded as random variables, i.e. we assume that $\mathbf{s}_1, \ldots, \mathbf{s}_n$ are independent and identically distributed random vectors. In this case, it is also usual to assume that they are independent of the measurement errors $\mathbf{e}_1, \ldots, \mathbf{e}_n$. Moreover, the normality assumption for the joint distribution of $\mathbf{s}_t$ and $\mathbf{e}_t$ is assumed. The parameters of the joint distribution of $\mathbf{w}_t$ and $\mathbf{s}_t$ is denoted by $\boldsymbol{\delta}$.

Let $(y_1, \mathbf{w}_1), \ldots, (y_n, \mathbf{w}_n)$ be the observable variables. We omit the observable vectors $\mathbf{z}_t$ and $\mathbf{v}_t$ in the notation as they are non-random and known. The joint density function of $(y_t, \mathbf{w}_t)$, which is the observation for the t-th individual, is obtained by integrating the joint density of the complete data $(y_t, \mathbf{w}_t, \mathrm{s}_t)$,

$$f(y_t, \mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\theta}, \boldsymbol{\delta}) = f(y_t|\mathbf{w_t}, \mathrm{s}_t; \boldsymbol{\theta}) f(\mathrm{s}_t, \mathbf{w}_t; \boldsymbol{\delta}),$$

with respect to s_t. Here, $\boldsymbol{\theta} = (\boldsymbol{\alpha}^\top, \boldsymbol{\beta}^\top, \boldsymbol{\gamma}^\top, \boldsymbol{\lambda}^\top)^\top$ represents the parameter of interest, and $\boldsymbol{\delta}$ is the nuisance parameter. The joint density $f(\mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\delta})$, which is associated to the measurement error model, can be written as $f(\mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\delta}) = f(\mathbf{w}_t|\mathrm{s}_t; \boldsymbol{\delta}) f(\mathrm{s}_t|\boldsymbol{\delta})$ as well as $f(\mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\delta}) = f(\mathrm{s}_t|\mathbf{w}_t; \boldsymbol{\delta}) f(\mathbf{w}_t|\boldsymbol{\delta})$. In this work we assume that, given the true (unobservable) covariates s_t, the response variable y_t does not depend on the surrogate covariates $\mathbf{w}_t$; i.e. $f(y_t|\mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\theta}) = f(y_t|\mathrm{s}_t; \boldsymbol{\theta})$. In other words, conditionally on s_t, y_t and $\mathbf{w}_t$ are assumed to be independent (Bolfarine and Arellano-Valle, 1998). Therefore, the density function of $(y_t, \mathbf{w}_t)$ is given by

$$\begin{aligned} f(y_t, \mathbf{w}_t; \boldsymbol{\theta}, \boldsymbol{\delta}) &= \int_{-\infty}^{\infty} \ldots \int_{-\infty}^{\infty} f(y_t, \mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\theta}, \boldsymbol{\delta}) d\mathrm{s}_t, \\ &= \int_{-\infty}^{\infty} \ldots \int_{-\infty}^{\infty} f(y_t|\mathrm{s}_t; \boldsymbol{\theta}) f(\mathbf{w}_t, \mathrm{s}_t; \boldsymbol{\delta}) d\mathrm{s}_t. \end{aligned}$$

The log-likelihood function for a sample of n observations is given by

$$\begin{aligned} \ell(\boldsymbol{\theta}, \boldsymbol{\delta}) &= \sum_{t=1}^{n} \log \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f(y_t|\mathrm{s}_t; \boldsymbol{\theta}) f(\mathrm{s}_t|\mathbf{w}_t; \boldsymbol{\delta}) f(\mathbf{w}_t; \boldsymbol{\delta}) d\mathrm{s}_t, \\ &= \sum_{t=1}^{n} \log f(\mathbf{w}_t; \boldsymbol{\delta}) + \sum_{i=1}^{n} \log \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f(y_t|\mathrm{s}_t; \boldsymbol{\theta}) f(\mathrm{s}_t|\mathbf{w}_t; \boldsymbol{\delta}) d\mathrm{s}_t. \end{aligned} \quad (7)$$

In general, the likelihood function involves analytically intractable integrals and, hence, approximate inference methods need to be considered. In the next section, we present three different approaches to estimate the parameters.

In order to facilitate the description of the estimation methods, we assume that a single covariate, x_t, is measured with error, and that it is used for the specifications of both the mean and precision submodels. We then have $p_\beta = p_\lambda = 1$ and $\mathrm{s}_t = x_t = m_t$. We also assume independence and normality of random errors. The methodologies presented in this paper can be extended to the situation where $\mathbf{x}_t$ and $\mathbf{m}_t$ are distinct, or when covariates measured with error appear only in the specification of the mean or the precision parameter. To be specific, from now on, the model under consideration is summarized as follows. For $i = 1, \dots, n$,

$$y_t | x_t, w_t \sim \mathrm{Beta}(\mu_t, \phi_t), \tag{8}$$

$$g(\mu_t) = \mathbf{z}_t^\top \boldsymbol{\alpha} + x_t\beta, \quad h(\phi_t) = \mathbf{v}_t^\top \boldsymbol{\gamma} + x_t\lambda, \tag{9}$$

$$w_t = \tau_0 + \tau_1 x_t + e_t, \;\; x_t \overset{\text{ind}}{\sim} N(\mu_x, \sigma_x^2), \;\; e_t \overset{\text{ind}}{\sim} N(0, \sigma_e^2), \tag{10}$$

with x_t and $e_{t'}$, for $t, t' = 1, \dots, n$, being independent. The unknown parameter vectors $\boldsymbol{\alpha}$ and $\boldsymbol{\gamma}$ were defined above, and $\beta \in \mathbb{R}$, $\lambda \in \mathbb{R}$, $\mu_x \in \mathbb{R}$ and $\sigma_x^2 > 0$ are unknown parameters. Note that it is assumed that the conditional distribution of y_t given (x_t, w_t) does not depend on w_t. Also, if $\tau_0 = 0$ and $\tau_1 = 1$, (10) corresponds to the classical additive error model $w_t = x_t + e_t$.

From (10) we have

$$w_t \overset{\text{ind}}{\sim} \mathrm{N}(\tau_0 + \tau_1\mu_x, \tau_1^2\sigma_x^2 + \sigma_e^2), \;\; x_t|w_t \overset{\text{ind}}{\sim} \mathrm{N}(\mu_{x_t|w_t}, \sigma_{x_t|w_t}^2), \tag{11}$$

where

$$\mu_{x_t|w_t} = \mu_x + k_x[w_t - (\tau_0 + \tau_1\mu_x)], \;\; \sigma_{x_t|w_t}^2 = \sigma_e^2 k_x/\tau_1, \tag{12}$$

with $k_x = \tau_1\sigma_x^2/(\tau_1^2\sigma_x^2 + \sigma_e^2)$ being known as the reliability ratio. To avoid non-identifiability of parameters we assume that $(\tau_0, \tau_1, \sigma_e^2)$ or (τ_0, τ_1, k_x) are either known parameters or they are estimated from supplementary information, typically replicate measurements or partial observation of the error-free covariate. In any case, either of these vectors are regarded as known quantities in the inferential procedure. Hence, the nuisance parameter vector is $\boldsymbol{\delta} = (\mu_x, \sigma_x^2)^\top$.

The log-likelihood function in (7) for n observations taken from the model described in (8), (9) and (10) is given by

$$\ell(\boldsymbol{\theta}, \boldsymbol{\delta}) = \sum_{t=1}^{n} \ell_{1t}(\boldsymbol{\delta}) + \sum_{t=1}^{n} \ell_{2t}(\boldsymbol{\theta}, \boldsymbol{\delta}), \tag{13}$$

where

$$
\begin{aligned}
\ell_{1t}(\boldsymbol{\delta}) &= -\frac{1}{2}\log[2\pi(\tau_1^2\sigma_x^2+\sigma_e^2)] - \frac{[w_t-(\tau_0+\tau_1\mu_x)]^2}{2(\tau_1^2\sigma_x^2+\sigma_e^2)}, \qquad (14)\\
\ell_{2t}(\boldsymbol{\theta},\boldsymbol{\delta}) &= \log\int_{-\infty}^{\infty} f(y_t|x_t;\boldsymbol{\theta},\boldsymbol{\delta})\frac{1}{\sqrt{2\pi\sigma^2_{x_t|w_t}}}\exp\left[-\frac{(x_t-\mu_{x_t|w_t})^2}{2\sigma^2_{x_t|w_t}}\right]dx_t. \qquad (15)
\end{aligned}
$$

3. Estimation

3.1. Maximum likelihood estimation

The second term of the log-likelihood function $\ell(\boldsymbol{\theta},\boldsymbol{\delta})$ in (13), which depends on a non-analytical integral (as can be seen in (15)), can be approximated using the Gauss-Hermite quadrature, which consists of the approximation

$$
\int_{-\infty}^{\infty} e^{-x^2} f(x)dx \approx \sum_{q=1}^{Q} \nu_q f(\eta_q), \qquad (16)
$$

where η_q and ν_q represent the q-th zero and weight, respectively, of the orthogonal Hermite polynomial of order Q (number of quadrature points); see, for instance, Abramowitz and Stegun (1972, Section 20.4.46). Using the transformation $u_t = (x_t-\mu_{x_t|w_t})/\sqrt{2\sigma^2_{x_t|w_t}}$ in (15), we have that $x_t = \mu_{x_t|w_t} + \sqrt{2\sigma^2_{x_t|w_t}}u_t$ and $dx_t = \sqrt{2\sigma^2_{x_t|w_t}}du_t$. Hence, by applying (16) in (15), the log-likelihood function (13) can be approximated by

$$
\ell_a(\boldsymbol{\theta},\boldsymbol{\delta}) = \sum_{t=1}^{n}\ell_{1t}(\boldsymbol{\delta}) + \sum_{t=1}^{n}\log\left(\sum_{q=1}^{Q}\frac{\nu_q}{\sqrt{\pi}}\exp[l(\mu_{tq},\phi_{tq})]\right),
$$

where

$$
\begin{aligned}
l(\mu,\phi) &= \log\Gamma(\phi) - \log\Gamma(\mu\phi) - \log\Gamma[(1-\mu)\phi] \qquad (17)\\
&+ (\mu\phi-1)\log(y_t) + [(1-\mu)\phi-1]\log(1-y_t),\\
g(\mu_{tq}) &= \mathbf{z}_t^\top\boldsymbol{\alpha} + x_t^*\beta,\;\; h(\phi_{tq}) = \mathbf{v}_t^\top\boldsymbol{\gamma} + x_t^*\lambda,\;\; x_t^* = \mu_{x_t|w_t} + \sqrt{2\sigma^2_{x_t|w_t}}\eta_q,
\end{aligned}
$$

where $\mu_{x_t|w_t}$ and $\sigma^2_{x_t|w_t}$ are given in (12).

For n and Q sufficiently large, the distribution of $(\widehat{\boldsymbol{\theta}}^\top, \widehat{\boldsymbol{\delta}}^\top)^\top$, the estimator of $(\boldsymbol{\theta}^\top, \boldsymbol{\delta}^\top)^\top$ obtained by solving the system of equations $\partial\ell_a(\boldsymbol{\theta}, \boldsymbol{\delta})/\partial\boldsymbol{\theta} = 0$, $\partial\ell_a(\boldsymbol{\theta}, \boldsymbol{\delta})/\partial\boldsymbol{\delta} = 0$ is approximately normally distributed with mean $(\boldsymbol{\theta}^\top, \boldsymbol{\delta}^\top)^\top$ and covariance matrix $\mathbf{J}_a^{-1}(\boldsymbol{\theta}, \boldsymbol{\delta})$, where

$$\mathbf{J}_a(\boldsymbol{\theta}, \boldsymbol{\delta}) = -\frac{\partial^2\ell(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial(\boldsymbol{\theta}^\top, \boldsymbol{\delta}^\top)^\top \partial(\boldsymbol{\theta}^\top, \boldsymbol{\delta}^\top)}$$

(Guolo, 2011). For computational implementation, the derivatives of $\ell_a(\boldsymbol{\theta}, \boldsymbol{\delta})$ with respect to the parameters can be analytically obtained or numerical derivatives can be used. Our numerical results were obtained using numerical derivatives.

3.2. Maximum pseudo-likelihood estimation

The central idea of the maximum pseudo-likelihood estimation method is to replace the nuisance parameters by consistent estimates in the log-likelihood function (13). The resulting function can be regarded as a pseudo-log-likelihood function that depends on the parameters of interest only (Guolo, 2011).

The log-likelihood function (13) is maximized in two steps. First, we estimate the nuisance parameter vector $\boldsymbol{\delta}$ by maximizing the reduced log-likelihood function

$$\ell_r(\boldsymbol{\delta}) = \sum_{t=1}^{n} \ell_{1t}(\boldsymbol{\delta}), \tag{18}$$

where $\ell_{1t}(\boldsymbol{\delta})$ is given in (14). Second, the estimate $\widehat{\boldsymbol{\delta}}$ obtained from the maximization of (18) is inserted in the original log-likelihood function (13), which results in the pseudo-log-likelihood function

$$\ell_p(\boldsymbol{\theta}; \widehat{\boldsymbol{\delta}}) = \sum_{t=1}^{n} \ell_{1t}(\widehat{\boldsymbol{\delta}}) + \sum_{t=1}^{n} \ell_{2t}(\boldsymbol{\theta}, \widehat{\boldsymbol{\delta}}). \tag{19}$$

As in $\ell(\boldsymbol{\theta}, \boldsymbol{\delta})$, the second term in $\ell_p(\boldsymbol{\theta}; \widehat{\boldsymbol{\delta}})$ is analytically intractable. However, unlike the integral in $\ell(\boldsymbol{\theta}, \boldsymbol{\delta})$, the integral in $\ell_p(\boldsymbol{\theta}; \widehat{\boldsymbol{\delta}})$ depends on the parameter of interest only. From (16), it is possible to approximate $\ell_{2t}(\boldsymbol{\theta}, \widehat{\boldsymbol{\delta}})$ by a summation. After some algebra, we have that an approximate pseudo-log-likelihood for the beta regression model with one covariate measured with error is given by (19) with $\ell_{2t}(\boldsymbol{\theta}, \widehat{\boldsymbol{\delta}})$ replaced by

$$\ell^{\dagger}_{2t}(\boldsymbol{\theta}, \widehat{\boldsymbol{\delta}}) \;=\; \log\left(\sum_{q=1}^{Q} \frac{\nu_q}{\sqrt{\pi}} \exp\left[l(\widehat{\mu}_{tq}, \widehat{\phi}_{tq})\right]\right),$$

where $l(\mu, \phi)$ is given in (17),

$$g(\widehat{\mu}_{tq}) = \mathbf{z}_t^{\top}\boldsymbol{\alpha} + \widehat{x}_t\beta, \;\; h(\widehat{\phi}_{tq}) = \mathbf{v}_t^{\top}\boldsymbol{\gamma} + \widehat{x}_t\lambda, \;\; \widehat{x}_t = \widehat{\mu}_{x_t|w_t} + \sqrt{2\widehat{\sigma}^2_{x_t|w_t}}\eta_q,$$

with $\widehat{\mu}_{x_t|w_t}$ and $\widehat{\sigma}^2_{x_t|w_t}$ being estimates of $\mu_{x_t|w_t}$ and $\sigma^2_{x_t|w_t}$, respectively, obtained from the maximization of (18).

It can be shown that, under regularity conditions (Gong and Samaniego, 1981), the asymptotic distribution of $\sqrt{n}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta})$ is normal with mean zero and covariance matrix

$$\boldsymbol{\Sigma} = \mathrm{I}_{\boldsymbol{\theta\theta}}^{-1}\left[\boldsymbol{\Sigma}_{\boldsymbol{\theta\theta}} - \mathrm{I}_{\boldsymbol{\theta\delta}}\mathrm{I}_{\boldsymbol{\delta\delta}}^{-1}\boldsymbol{\Sigma}_{\boldsymbol{\delta\theta}} - \boldsymbol{\Sigma}_{\boldsymbol{\delta\theta}}^{\top}\mathrm{I}_{\boldsymbol{\delta\delta}}^{-1}\mathrm{I}_{\boldsymbol{\theta\delta}}^{\top} + \mathrm{I}_{\boldsymbol{\theta\delta}}\mathrm{I}_{\boldsymbol{\delta\delta}}^{-1}\boldsymbol{\Sigma}_{\boldsymbol{\delta\delta}}\mathrm{I}_{\boldsymbol{\delta\delta}}^{-1}\mathrm{I}_{\boldsymbol{\theta\delta}}^{\top}\right]\mathrm{I}_{\boldsymbol{\theta\theta}}^{-1},$$

where

$$\begin{aligned}
\mathrm{I}_{\boldsymbol{\theta\theta}} &= -\sum_{t=1}^{n}\mathrm{E}\left(\frac{\partial^2\ell_{pt}(\boldsymbol{\theta},\boldsymbol{\delta})}{\partial\boldsymbol{\theta}\partial\boldsymbol{\theta}^{\top}}\right), \;\; \mathrm{I}_{\boldsymbol{\theta\delta}} = -\sum_{t=1}^{n}\mathrm{E}\left(\frac{\partial^2\ell_{pt}(\boldsymbol{\theta},\boldsymbol{\delta})}{\partial\boldsymbol{\theta}\partial\boldsymbol{\delta}^{\top}}\right),\\
\boldsymbol{\Sigma}_{\boldsymbol{\theta\theta}} &= \sum_{t=1}^{n}\frac{\partial\ell_{pt}(\boldsymbol{\theta},\boldsymbol{\delta})}{\partial\boldsymbol{\theta}}\left(\frac{\partial\ell_{pt}(\boldsymbol{\theta},\boldsymbol{\delta})}{\partial\boldsymbol{\theta}}\right)^{\top}, \;\; \boldsymbol{\Sigma}_{\boldsymbol{\delta\theta}} = \sum_{t=1}^{n}\frac{\partial\ell_{rt}(\boldsymbol{\delta})}{\partial\boldsymbol{\delta}}\left(\frac{\partial\ell_{pt}(\boldsymbol{\theta},\boldsymbol{\delta})}{\partial\boldsymbol{\theta}}\right)^{\top},\\
\boldsymbol{\Sigma}_{\boldsymbol{\delta\delta}} &= \sum_{t=1}^{n}\frac{\partial\ell_{rt}(\boldsymbol{\delta})}{\partial\boldsymbol{\delta}}\left(\frac{\partial\ell_{rt}(\boldsymbol{\delta})}{\partial\boldsymbol{\delta}}\right)^{\top},
\end{aligned}$$

with $\boldsymbol{\delta}$ replaced by $\widehat{\boldsymbol{\delta}}$, and with $\ell_{rt}(\boldsymbol{\delta})$ and $\ell_{pt}(\boldsymbol{\theta}, \boldsymbol{\delta})$ being the t-th element of the log-likelihood functions $\ell_r(\boldsymbol{\delta})$ and $\ell_p(\boldsymbol{\theta}, \boldsymbol{\delta})$ in (18) and (19), respectively (Guolo, 2011). Further details can be found in Carroll *et al.* (2006, Section A.6.6). For the errors-in-variables beta regression model considered here, $\mathrm{I}_{\boldsymbol{\theta\theta}}$, $\mathrm{I}_{\boldsymbol{\delta\delta}}$ and $\mathrm{I}_{\boldsymbol{\theta\delta}}$ do not have closed form. We suggest to replace the expected information matrix by the observed information matrix. For computational implementation, the needed derivatives can be analytically or numerically obtained. We used numerical derivatives in our applications.

3.3. Regression calibration estimation

The regression calibration method has been widely used in errors-in-variables modeling due to its simplicity; see Carroll *et al.* (2006, Chap. 4), Freedman *et al.* (2008), Thurston *et al.* (2005) and Guolo (2011). The central idea is to replace the unobservable variable, x_t, by an estimate of the conditional expected value of x_t given w_t, $\mathrm{E}(x_t|w_t)$, in the likelihood function. Let $r(w_t, \boldsymbol{\delta}) = \mathrm{E}(x_t|w_t)$ be the calibration function. The replacement of the unobservable covariate x_t by $r(w_t, \widehat{\boldsymbol{\delta}})$ establishes a modified model for the data. Here, $\widehat{\boldsymbol{\delta}}$ is an estimate of $\boldsymbol{\delta}$.

For our errors-in-variables beta regression model, the calibration function is $r(w_t, \boldsymbol{\delta}) = \mu_{x_t|w_t}$ as defined in (12) . From (11) and (12), we have that $\overline{w} = \sum_{t=1}^{n} w_t/n$ and $s_w^2 = \sum_{t=1}^{n}(w_t - \overline{w})^2/(n-1)$ are optimal estimates of $\tau_0 + \tau_1\mu_x$ and $\tau_1^2\sigma_x^2 + \sigma_e^2$, respectively. These estimates can be used to estimate the calibration function.

By inserting the calibration function in the conditional density function of y_t given x_t, we obtain the modified log-likelihood function

$$\ell_{rc}(\boldsymbol{\theta}) = \sum_{t=1}^{n} l(\widetilde{\mu}_t, \widetilde{\phi}_t), \tag{20}$$

where $l(\mu, \phi)$ is given in (17),

$$g(\widetilde{\mu}_t) = \mathbf{z}_t^\top \boldsymbol{\alpha} + \widetilde{x}_t\beta, \quad h(\widetilde{\phi}_t) = \mathbf{v}_t^\top \boldsymbol{\gamma} + \widetilde{x}_t\lambda,$$

with $\widetilde{x}_t$ being the estimated calibration function. Note that the modified log-likelihood function in (20) only depends on the parameter of interest, $\boldsymbol{\theta}$. The regression calibration estimate of $\boldsymbol{\theta}$ is obtained from the system of equations $\partial\ell_{rc}(\boldsymbol{\theta})/\partial\boldsymbol{\theta} = 0$, which requires a numerical algorithm. Since $\ell_{rc}(\boldsymbol{\theta})$ coincides with the log-likelihood function for the usual beta regression model, $\widetilde{x}_t$ acting as an observable covariate, these equations can be numerically solved from available computational packages, for instance the `betareg` package (Cribari-Neto and Zeiles, 2010) implemented in the `R` platform.

We note that (20) is a pseudo-likelihood as presented in Gong and Samaniego (1981). Gong and Samaniego (1981) present the maximum pseudo-likelihood method as follows. Consider a sample of size n with joint density function $f(\cdot; \boldsymbol{\theta}, \boldsymbol{\delta})$, and assume that an estimate $\widehat{\boldsymbol{\delta}}$ of $\boldsymbol{\delta}$ is available. Hence, the maximum pseudo-likelihood estimate, $\widehat{\boldsymbol{\theta}}$, is obtained by maximizing $f(\cdot; \boldsymbol{\theta}, \widehat{\boldsymbol{\delta}})$. One expects $\widehat{\boldsymbol{\theta}}$ to have good properties for large samples whenever $\widehat{\boldsymbol{\delta}}$ has; for further details see

Delgado (1995). From Gong and Samaniego (1981), Parke (1986) and Delgado (1995) it can be shown that, under regularity conditions, the asymptotic distribution of $\sqrt{n}(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta})$ is normal with mean zero and covariance matrix

$$\boldsymbol{\Sigma}^* = \mathrm{I}^{*-1}_{\boldsymbol{\theta\theta}} \left(\mathrm{I}^*_{\boldsymbol{\theta\theta}} + \mathrm{I}^*_{\boldsymbol{\theta\delta}} \boldsymbol{\Sigma}^*_{\boldsymbol{\delta\delta}} \mathrm{I}^{*\top}_{\boldsymbol{\theta\delta}} \right) \mathrm{I}^{*-1}_{\boldsymbol{\theta\theta}},$$

where

$$\begin{aligned}
\mathrm{I}^*_{\boldsymbol{\theta\theta}} &= \sum_{t=1}^{n} \mathrm{E}\left[\frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\theta}} \left(\frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\theta}} \right)^{\top} \right], \\
\mathrm{I}^*_{\boldsymbol{\theta\delta}} &= \sum_{t=1}^{n} \mathrm{E}\left[\frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\theta}} \left(\frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\delta}} \right)^{\top} \right], \\
\boldsymbol{\Sigma}^*_{\boldsymbol{\delta\delta}} &= \sum_{t=1}^{n} \frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\delta}} \left(\frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\delta}} \right)^{\top},
\end{aligned}$$

with $\boldsymbol{\delta}$ replaced by $\widehat{\boldsymbol{\delta}}$ and with $\ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})$ being the t-th term of $\ell_{rc}(\boldsymbol{\theta}, \boldsymbol{\delta})$. To obtain $\boldsymbol{\Sigma}^*$ one needs to obtain $\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})/\partial \boldsymbol{\delta}$, $\mathrm{I}^*_{\boldsymbol{\theta\theta}}$ and $\mathrm{I}^*_{\boldsymbol{\theta\delta}}$. We have

$$\frac{\partial \ell_{rct}(\boldsymbol{\theta}, \boldsymbol{\delta})}{\partial \boldsymbol{\delta}} = \left[\begin{array}{r} \beta(1 - \tau_1 k_x)\phi_t(y_t^* - \mu_t^*)g'(\mu_t)^{-1} + \lambda(1 - \tau_1 k_x)a_t h'(\phi_t)^{-1} \\ \beta(1 - \tau_1 k_x)\phi_t(y_t^* - \mu_t^*)g'(\mu_t)^{-1}\sigma_x^{-2} + \lambda(1 - \tau_1 k_x)a_t h'(\phi_t)^{-1}\sigma_x^{-2} \end{array} \right],$$

where

$$a_t = \mu_t(y_t^* - \mu_t^*) + \log(1 - y_t) - \psi((1 - \mu_t)\phi_t) + \psi(\phi_t),$$

$$y_t^* = \log\left(\frac{y_t}{1 - y_t} \right), \quad \mu^* = \psi(\mu_t \phi) - \psi[(1 - \mu_t)\phi] \tag{21}$$

with $\psi(\cdot)$ being the digamma function, i.e., $\psi(z) = \mathrm{d}\log\Gamma(z)/\mathrm{d}z$ for $z > 0$. Here, we are able to obtain $\mathrm{I}^*_{\boldsymbol{\theta\theta}}$ and $\mathrm{I}^*_{\boldsymbol{\theta\delta}}$, since the log-likelihood function does not involve integrals. After some algebra, we have

$$\mathrm{I}^*_{\boldsymbol{\theta\theta}} = \left[\begin{array}{cccc} \mathbf{Z}^\top \Phi \mathbf{U} \mathbf{Z} & \mathbf{Z}^\top \Phi \mathbf{U} \mathbf{x} & \mathbf{Z}^\top \mathbf{C} \mathbf{G} \mathbf{H} \mathbf{V} & \mathbf{Z}^\top \mathbf{C} \mathbf{G} \mathbf{H} \mathbf{x} \\ & \mathbf{x}^\top \Phi \mathbf{U} \mathbf{x} & \mathbf{x}^\top \mathbf{C} \mathbf{G} \mathbf{H} \mathbf{V} & \mathbf{x}^\top \mathbf{C} \mathbf{G} \mathbf{H} \mathbf{x} \\ & & \mathbf{V}^\top \mathbf{D} \mathbf{V} & \mathbf{V}^\top \mathbf{D} \mathbf{x} \\ & & & \mathbf{x}^\top \mathbf{D} \mathbf{x} \end{array} \right],$$

where

$$\mathbf{Z} = (\mathbf{z}_1, \ldots, \mathbf{z}_n), \;\; \mathbf{V} = (\mathbf{v}_1, \ldots, \mathbf{v}_n), \;\; \mathbf{U} = \text{diag}\{u_1, \ldots, u_n\},$$

$$\Phi = \text{diag}\{\phi_1, \ldots, \phi_n\}, \;\; \mathbf{H} = \text{diag}\{h'(\phi_1)^{-1}, \ldots, h'(\phi_n)^{-1}\},$$
$$\mathbf{G} = \text{diag}\{g'(\mu_1)^{-1}, \ldots, g'(\mu_n)^{-1}\}, \;\; \mathbf{C} = \text{diag}\{c_1, \ldots, c_n\},$$
$$\mathbf{D} = \text{diag}\{d_1, \ldots, d_n\},$$

with

$$u_t = \phi_t \frac{\psi'(\mu_t\phi_t) + \psi'((1-\mu_t)\phi_t)}{g'(\mu_t)^2},$$
$$c_t = \phi_t[\psi'(\mu_t\phi_t)\mu_t - \psi'((1-\mu_t)\phi_t)(1-\mu_t)],$$
$$d_t = \frac{\psi'(\mu_t\phi_t)\mu_t^2 + \psi'((1-\mu_t)\phi_t)(1-\mu_t^2) - \psi'(\phi_t)}{h'(\phi_t)^2}.$$

Analogously, we have

$$\mathrm{I}^*_{\boldsymbol{\theta\delta}} = \begin{bmatrix} \mathbf{Z}^\top(\mathbf{U}^* + \mathbf{HGC}^*)\mathbf{1}_n & \mathbf{Z}^\top(\mathbf{U}^\dagger + \mathbf{HGC}^\dagger)\mathbf{1}_n \\ \mathbf{x}^\top(\mathbf{U}^* + \mathbf{HGC}^*)\mathbf{1}_n & \mathbf{x}^\top(\mathbf{U}^\dagger + \mathbf{HGC}^\dagger)\mathbf{1}_n \\ \mathbf{V}^\top(\mathbf{D}^* + \mathbf{HGC}^*)\mathbf{1}_n & \mathbf{V}^\top(\mathbf{D}^\dagger + \mathbf{HGC}^\dagger)\mathbf{1}_n \\ \mathbf{x}^\top(\mathbf{D}^* + \mathbf{HGC}^*)\mathbf{1}_n & \mathbf{x}^\top(\mathbf{D}^\dagger + \mathbf{HGC}^\dagger)\mathbf{1}_n \end{bmatrix},$$

where

$$\mathbf{U}^* = \text{diag}\{u_1^*, \ldots, u_n^*\}, \;\; \mathbf{U}^\dagger = \text{diag}\{u_1^\dagger, \ldots, u_n^\dagger\}, \;\; \mathbf{C}^* = \text{diag}\{c_1^*, \ldots, c_n^*\},$$

$$\mathbf{C}^\dagger = \text{diag}\{c_1^\dagger, \ldots, c_n^\dagger\}, \;\; \mathbf{D}^* = \text{diag}\{d_1^*, \ldots, d_n^*\}, \mathbf{D}^\dagger = \text{diag}\{d_1^\dagger, \ldots, d_n^\dagger\},$$

with

$$u_t^* = \beta(1-\tau_1 k_x)u_t, \;\; u_t^\dagger = u_t^* k_x \sigma_x^{-2}, \;\; c_t^* = \beta(1-\tau_1 k_x)c_t,$$
$$c_t^\dagger = c^* k_x \sigma_x^{-2}, \;\; d_t^* = \lambda(1-\tau_1 k_x)d_t, \;\; d_t^\dagger = d_t^* k_x \sigma_x^{-2},$$

and $\mathbf{1}_n$ is a vector of ones($n \times 1$).

4. Monte Carlo simulation results

We now present Monte Carlo simulation results on the performance of the different estimation methods described in Section 3. All simulation results are based on 5,000 Monte Carlo replications. We consider errors-in-variables beta regression models with $\log(\mu_t/(1-\mu_t)) = \alpha + \beta x_t$, and $\log(\phi_t) = \gamma$ (constant precision model) and $\log(\phi_t) = \gamma + \lambda x_t$ (varying precision model), with w_t and x_t being generated from (10). We set α =2.0, $\beta = -0.6$, $\lambda = 0.5$, $\mu_x = 2.5$, $\sigma_x^2 = 2.7$, and $\gamma = 2.5$ for the constant precision model and $\gamma = 4$ for the varying precision model. The parameters of the measurement error mechanism are assumed to be known, and we set $\tau_0 = 0$, $\tau_1 = 1$, and the following values for σ_e^2: 0.05 and 0.50 ($k_x = 0.98$, and 0.84, respectively). The sample sizes are $n = 25$, 50, 100, 200, and 300. For each simulated sample the parameters were estimated under two different settings. First, we ignored the measurement error in x_t, i.e. we used what is called the naïve method (ℓ_{naive}); second, we recognized that x_t is measured with error and estimated the parameters using the three methods proposed in this paper: approximate maximum likelihood (ℓ_a), (approximate) maximum pseudo-likelihood (ℓ_p), and regression calibration (ℓ_{rc}). Whenever Gaussian quadrature was required, use set the number of quadrature points at $Q = 50$. The maximization of the relevant (approximate/pseudo/modified) likelihoods was performed using the quasi-Newton BFGS nonlinear optimization algorithm with numerical derivatives implemented in the function MAXBFGS in the matrix language programming `Ox` (Doornik, 2011).

Figures 1-4 present plots of the bias and the root-mean-square error of the estimators against sample size, for $k_x = 0.98$ and 0.84 under the constant precision model and the varying precision model. As expected, the naïve estimator is biased and its bias and mean-square error do not converge to zero as n grows even when the reliability ratio is large (i.e. the variance of the measurement error is small compared to the variance of x). In other words, the plots suggest that the naïve estimator is not consistent. For all the cases, the approximate maximum likelihood and maximum pseudo-likelihood estimators perform similarly. In general, their performance is clearly better than that of the regression calibration and naïve estimators. Under constant precision (Figures 1-2), the regression calibration estimator is as biased as the naïve estimator for estimating the precision parameter. However, for estimating β, the coefficient associated to the covariate measured with error, it performs well if the measurement error variance is small. In fact, the regression calibration, approximate maximum likelihood ratio and maximum pseudo-likelihood ratio estimators are virtually unbiased for estimating β when $k_x = 0.98$; also, their mean-square errors converge to zero as n grows. Under the varying precision model (Figures 3-4), similar conclusions are reached.

We now turn to the evaluation of confidence intervals constructed from the different estimators coupled with their respective asymptotic distributions. Figures 5-6 present plots of the estimated true coverages of confidence intervals constructed with 95% nominal confidence level, for $k_x = 0.98$ and 0.84 under the constant precision and varying precision models for n ranging from 25 to 300. For all the cases, the estimated true coverages of the confidence intervals based on the naïve estimator decrease as n grows, and hence they cannot be recommended. For all the cases, the confidence intervals constructed from the approximate maximum likelihood and maximum pseudo-likelihood estimators present true coverage close to 95%, more so if the sample size is large. For the constant precision model (Figure 5), the regression calibration estimator produces reliable confidence intervals for parameters of the mean submodel when the measurement error variance is small. However, for estimating the precision parameter, the regression calibration estimator produces confidence intervals with true coverage much smaller than 95% when n is large. Under the varying precision model (Figure 6), we arrive at similar conclusions.

Overall, we conclude that ignoring the measurement error produces misleading inference. Also, if the measurement error variance is small, the regression calibration approach is reliable for estimating the parameters of the mean submodel. Moreover, inference based on the approximate likelihood and the pseudo-likelihood methods present good performance for the estimation of all the parameters. Since the pseudo-likelihood approach is computationally less demanding than the approximate maximum likelihood approach, we recommend the maximum pseudo-likelihood estimation for practical applications.

5. Residual analysis

Espinheira *et al.* (2008) and Ferrari *et al.* (2011) proposed the use of standardized weighted residuals as a diagnostic tool for beta regression models with constant and non-constant precision parameter, respectively. Here, we modify the residuals defined by Ferrari *et al.* (2011) to allow for measurement error in covariates. We therefore define

$$r_t = \frac{y_t^* - \widehat{\mu}_t^*}{\sqrt{\widehat{\upsilon}_t(1 - \widehat{h}_{tt}^*)}},$$

for $t = 1, \dots, n$, where y_t^* and μ_t^* are given in (21), $\upsilon_t = \psi'(\mu_t\phi_t) + \psi'[(1-\mu_t)\phi_t]$ and h_{tt}^* is the t-th diagonal element of

$$\mathbf{H}^* = (\mathbf{M\Phi})^{1/2}\mathbf{W}(\mathbf{W}^\top\mathbf{\Phi M W})^{-1}\mathbf{W}^\top(\mathbf{\Phi M})^{1/2},$$



Figure 1: Bias and RMSE for the estimators of α, β and γ for $k_x = 0.98$, constant precision model; ℓ_a (square), ℓ_p (circle), ℓ_{rc} (triangle) and ℓ_{naive}(star).



Figure 2: Bias and RMSE for the estimators of α, β and γ for $k_x = 0.84$, constant precision model; ℓ_a (square), ℓ_p (circle), ℓ_{rc} (triangle) and ℓ_{naive}(star).



Figure 3: Bias and RMSE for the estimators of α, β, γ and λ for $k_x = 0.98$, varying precision model; ℓ_a (square), ℓ_p (circle), ℓ_{rc} (triangle) and ℓ_{naive}(star).



Figure 4: Bias and RMSE for the estimators of α, β, γ and λ for $k_x = 0.84$, varying precision model; ℓ_a (square), ℓ_p (circle), ℓ_{rc} (triangle) and ℓ_{naive}(star).



(a) (b)

Figure 5: Coverage of confidence intervals of α, β and γ for: column (a) $k_x = 0.98$, constant precision model, and column (b) $k_x = 0.84$, constant precision model; ℓ_a (square), ℓ_p (circle), ℓ_{rc} (triangle) and ℓ_{naive}(star).



(a) (b)

Figure 6: Coverage of confidence intervals of α, β, γ and λ for: column (a) $k_x = 0.98$, varying precision model, and column (b) $k_x = 0.84$, varying precision model; ℓ_a (square), ℓ_p (circle), ℓ_{rc} (triangle) and ℓ_{naive}(star).

in which $\mathbf{M} = \operatorname{diag}\{m_1, \ldots, m_n\}$ with $m_t = \phi_t v_t/[g^{'}(\mu_t)]^2$, $\boldsymbol{\Phi} = \operatorname{diag}\{\phi_1, \ldots, \phi_n\}$ and $\mathbf{W}$ is an $n \times (p_\alpha + p_\beta)$ matrix with the t-th row given by $(\mathbf{z}_t^\top, \mathbf{w}_t^\top)$. Here, hat indicates that the unknown parameters are replaced by estimates. We suggest the use of the maximum pseudo-likelihood estimates since they performed well in our simulations and are computationally less demanding than the approximate maximum likelihood estimates.

Plots of residuals versus observation indices are not always suitable for detecting lack of fit when measurement errors are present; see, for instance, Fuller (1987, Section 2.2.2), Carroll and Spiegelman (1992) and Buonaccorsi (2010, Section 4.7). Fuller (1987, Section 2.2.2) suggests constructing plots of residuals versus consistent estimates of the expected value of x_t given w_t. Atkinson (1985), on the other hand, suggests the use of simulated envelopes in normal probability plots to facilitate their interpretation. The use of standardized weighted residuals plots proposed here will be illustrated in the next section.

6. Real data application

We now illustrate our results in the dataset described in Section 1. Our aim is to model RPP (relative pulse pressure) using BMI (body mass index) and LDL (low density lipoprotein cholesterol level) as covariates. The total cholesterol level, TC, is easier to be accessed than LDL, and provides a measure of LDL plus unknown quantities of other components as triglycerides and high density lipoprotein. Here, we shall consider TC as a surrogate for LDL.

We shall assume that $y_1, y_2, \ldots, y_n$ are independent observations of the PPR of the n individuals in the sample, and such that y_t follows a beta distribution with mean μ_t and precision parameter ϕ_t, with

$$\begin{aligned}
\log[\mu_t/(1-\mu_t)] &= \alpha_0 + \alpha_1 \mathrm{BMI}_t + \beta \mathrm{LDL}_t, \\
\log(\phi_t) &= \gamma_0 + \gamma_1 \mathrm{BMI}_t + \lambda \mathrm{LDL}_t, \\
\mathrm{TC}_t &= \tau_0 + \tau_1 \mathrm{LDL}_t + e_t, \\
LDL_t &\overset{\text{ind}}{\sim} N(\mu_x, \sigma_x^2), \quad e_t \overset{\text{ind}}{\sim} N(0, \sigma_e^2),
\end{aligned} \tag{22}$$

for $t = 1, \ldots, 182$. Also, LDL_t and $e_{t'}$, for $t, t' = 1, \ldots, n$ are assumed to be independent Here, the interest parameter vector is $\boldsymbol{\theta} = (\alpha_0, \alpha_1, \beta, \gamma_0, \gamma_1, \lambda)^\top$.

For this particular dataset, the values of LDL and TC are available for all the individuals in the sample. We then use these data to mimic a situation where both the true and surrogate covariates are observed for some but not all individuals in the sample. We randomly selected a subsample of 21 individuals for which we consider the corresponding observed values of LDL and TC; for the remaining individuals, only the observations on TC are considered in the analysis. A scatter plot of TC versus LDL for the selected individuals (not shown) suggests a clear approximate linear tendency.

It is possible to estimate τ_0, τ_1 and σ_e^2 from (22), since we have observations on both TC and LDL for some individuals. We obtained the following estimates: $\widehat{\tau}_0 = 0.7351$, $\widehat{\tau}_1 = 1.062$ and $\widehat{\sigma}_e^2 = 0.030$. To estimate the parameters of interest we used the approximate maximum likelihood, maximum pseudo-likelihood, regression calibration and naïve methods. Also, as a gold standard to compare with these methods, we fitted a beta regression model in which LDL is used as the true covariate, measured without error, for all individuals.

Table 1 shows the estimates, standard errors and p-values for the parameters of interest. All the approaches produce similar inferences on the parameters of the mean submodel (α_0, α_1 and β) and on the intercept (γ_0) and the coefficient γ_1 of BMI, which is the covariate measured without error, in the precision submodel. Inference on λ, the coefficient of LDL (the covariate measured with error) in the precision submodel, varies depending on the approach being used. The gold standard and the approximate maximum likelihood and pseudo-likelihood approaches indicate that the null hypothesis $\mathcal{H}_0 : \lambda = 0$ should be rejected at the 5% nominal level (p-value$= 0.013,\ 0.025,\ 0.022$, respectively), while $\mathcal{H}_0$ is not rejected when the regression calibration and the naïve methods are employed (p-value$= 0.064,\ 0.064$). In other words, at the 5% nominal level, the approximate maximum likelihood and maximum pseudo-likelihood approaches agree with the gold standard in that they indicate that the precision varies with LDL, unlike the regression calibration and naïve methods. Finally, it can be noticed that the results for the approximate maximum likelihood and maximum pseudo-likelihood methods are very close.

We now use the standardized weighted residual presented in Section 5 to investigate the presence of outliers or any indication of lack of fit. Figure 7 shows residual plots for the model fitted using the maximum pseudo-likelihood approach. Figure 7(a) shows the plot of the residuals against predicted values of LDL, $\widehat{LDL}$, and Figure 7(b) shows a normal probability plot with simulated envelope. There is no indication of outliers or any apparent pattern. This indicates that the errors-in-variables model considered here fits the data well.

Table 1: Estimates, standard errors and p-values

Method	Parameter	Estimate	Standard error	z stat	p-value
Gold standard	α_0	−0.354	0.124	−2.855	0.004
	α_1	−0.009	0.004	−2.250	0.023
	β	0.107	0.050	2.140	0.034
	γ_0	5.905	0.764	7.729	0.000
	γ_1	−0.022	0.024	−0.916	0.356
	λ	−0.729	0.292	−2.497	0.013
ℓ_a	α_0	−0.366	0.113	−3.239	0.006
	α_1	−0.009	0.004	−2.250	0.024
	β	0.118	0.062	1.903	0.058
	γ_0	6.109	0.908	6.728	0.000
	γ_1	−0.028	0.028	−1.000	0.310
	λ	−0.751	0.336	−2.235	0.025
ℓ_p	α_0	−0.366	0.133	−2.759	0.006
	α_1	−0.009	0.004	−2.250	0.013
	β	0.118	0.068	1.735	0.082
	γ_0	6.109	0.994	6.146	0.000
	γ_1	−0.028	0.036	−0.778	0.434
	λ	−0.751	0.329	−2.283	0.022
ℓ_{rc}	α_0	−0.356	0.134	−2.657	0.008
	α_1	−0.009	0.004	−2.250	0.023
	β	0.112	0.062	1.806	0.068
	γ_0	5.945	0.804	7.394	0.000
	γ_1	−0.028	0.024	−1.167	0.239
	λ	−0.649	0.351	−1.849	0.064
ℓ_{naive}	α_0	−0.389	0.144	−2.701	0.007
	α_1	−0.009	0.004	−2.250	0.023
	β	0.085	0.047	1.809	0.068
	γ_0	6.136	0.864	7.102	0.000
	γ_1	−0.028	0.024	−1.167	0.239
	λ	−0.490	0.265	−1.849	0.064

7. Concluding remarks

In this paper we proposed and studied errors-in-variables beta regression models. We proposed three different estimation methods, namely, the approximate maximum likelihood, maximum pseudo-likelihood and regression calibration methods. We performed a Monte Carlo simulation study to compare the perfor-



(a) (b)

Figure 7: Plot of standardized weighted residuals versus predicted values of LDL (a), and normal probability plot of standardized weighted residuals (b).

mance of the estimators in terms of bias, root-mean-square errors and coverage of confidence intervals. Overall, we reached the following conclusions. First, ignoring the measurement error may lead to severely biased inference. Second, the regression calibration approach is very simple and is reliable for estimating the parameters of the mean submodel when the measurement error variance is small. Third, the approximate maximum likelihood and maximum pseudo-likelihood approaches perform well, the later being less computationally demanding than the former. We, therefore, recommend the maximum pseudo-likelihood estimation for practical applications. We also proposed a standardized weighted residual for diagnostic purposes. All our results were illustrated in the analysis of a real data set.

Acknowledgements

We gratefully acknowledge the financial support from CAPES-Brazil, CNPq-Brazil and FONDECYT(1120121)-Chile.

References

Abramowitz, M. and Stegun, I. A. (1972). *Handbook of Mathematical Functions*. New York:Dover.

American College of Surgeons (2008). *ATLS Advanced Trauma Life Support Program for Doctors*. Chicago: American College of Surgeons.

Arellano-Valle, R. B. and Bolfarine, H. (1996). A note on the simple structural regression model. *Annals of the Institute of Statistical Mathematics*, **48**, 111–125.

Atkinson, A. C. (1985). *Plots, Transformations and Regression: An Introduction to Graphical Methods of Diagnostic Regression Analysis*. New York: Oxford University Press.

Bolfarine, H. and Arellano-Valle, R. B. (1998). Weak nondifferential measurement error models. *Statistics and Probability Letters*, **40**, 279–287.

Buonaccorsi, J. P. (2010). *Measurement Error: Models, Methods and Applications*. London: Chapman and Hall.

Carroll, R. J. and Spiegelman, C. H. (1992). Diagnostics for nonlinearity and heteroscedasticity in errors-in-variables regression. *Technometrics*, **34**, 186–196.

Carroll, R. J., Ruppert, D., Stefanski, L. A., and Crainiceanu, C. M. (2006). *Measurement Error in Nonlinear Models: A Modern Perspective*. New York: Chapman and Hall.

Castro, M., Galea, M., and Bolfarine, H. (2008). Hypothesis testing in an errors-in-variables model with heteroscedastic measurement errors. *Statistics in Medicine*, **27**, 5217–5234.

Cheng, C. and Van Ness, J. W. (1999). *Statistical Regression with Measurement Error*. London: Oxford University Press.

Cribari-Neto, F. and Zeiles, A. (2010). Beta regression in R. *Journal of Statistical Software*, **34**, 1–24.

Delgado, J. J. F. (1995). *Pseudo-Maximum Likelihood Estimation*. Master's Thesis, University of São Paulo, in portuguese.

Doornik, J. (2011). *Ox 6.2: Object-Oriented Matrix Programming Language*. London: Timberlake Consultants and Oxford: http://www.doornik.com.

Espinheira, P. L., Ferrari, S. L. P., and Cribari-Neto, F. (2008). On beta regression residuals. *Journal of Applied Statistics*, **35**, 407–419.

Ferrari, S. L. P. and Cribari-Neto, F. (2004). Beta regression for modelling rates and proportions. *Journal of Applied Statistics*, **31**, 799–815.

Ferrari, S. L. P., Espinheira, P. L., and Cribari-Neto, F. (2011). Diagnostic tools in beta regression with varying dispersion. *Statistica Neerlandica*, **65**, 337–351.

Freedman, L. S., Midthune, D., Carroll, R., and Kipnis, V. (2008). A comparison of regression calibration, moment reconstruction and imputation for adjusting for covariate measurement error in regression. *Statistics in Medicine*, **27**, 5195–5216.

Fuller, W. A. (1987). *Measurement Error Models*. New York: John Wiley.

Gong, G. and Samaniego, F. J. (1981). Pseudo maximum likelihood estimation: theory and applications. *The Annals of Statistics*, **9**, 861–869.

Guolo, A. (2011). Pseudo-likelihood inference for regression models with misclassified and mismeasured variables. *Statistica Sinica*, **21**, 1639–1663.

Parke, W. R. (1986). Pseudo maximum likelihood estimation: the asymptotic distribution. *The Annals of Statistics*, **14**, 335–357.

Simas, A. B., Barreto-Souza, W., and Rocha, A. V. (2010). Improved estimators for a general class of beta regression models. *Computational Statistics and Data Analysis*, **54**, 348–366.

Smithson, M. and Verkuilen, J. (2006). A better lemon-squeezer? maximum-likelihood regression with beta-distributed dependent variables. *Psychological Methods*, **11**, 54–71.

Thurston, S. W., Williams, P. L., Hauser, R., Hu, H., Hernandez-Avila, M., and Spiegelman, D. (2005). A comparison of regression calibration approaches for designs with internal validation data. *Journal of Statistical Planning and Inference*, **131**, 175–190.